


ANEXOS

Tabla n°1. Riesgo relativo de fractura de cadera según factores de riesgo del adulto mayor.

Characteristics	Hazard ratio	95 % CI
Parkinson's disease	5.12	1.82–14.44
Currently smoking	4.38	2.06–9.33
Delirium in previous month	2.01	1.15–3.49
Underweight (BMI <22.0)	1.74	1.09–2.77
Age	1.09	1.04–1.14
Hip prosthesis	0.37	0.15–0.91
Female sex	0.875	0.56–1.37


Fuente: Wiklund R et al. Risk factors for hip fracture in very old people: a population-based study. *Osteoporos Int.* 2016; 27:923–931.

Grafico n°1. Riesgo relativo de fractura de cadera por valor de 25OHD sérico, ajustado por edad, sexo, raza/etnia.



Fuente: Looker A. Serum 25-hydroxyvitamin d and risk of major osteoporotic fractures in older U.S. adults. *Journal of Bone and Mineral Research.* 2013; 28(5): 997–1006.

Grafico n°2. Riesgo relativo de fractura de cadera por decil de hemoglobina.



Fuente: Looker A. Hemoglobin and hip fracture risk in older non-Hispanic white adults. *Osteoporos Int.* 2014; 25(10): 2389–2398.